

GUIDELINES FOR SURVEILLANCE AND RISK ANALYSIS OF PLANT PESTS IN NEPAL

Endorsed by National Plant Protection Organization-Nepal, on 2071/01/22 BS,
First updated on 2081/02/29, Second updated on 2082/02/20



**Government of Nepal
Ministry of Agriculture and Livestock Development
Plant Quarantine and Pesticide Management Center
Hariharbhawan, Lalitpur**

Tel.: +977-1-5524352, 5553798 Fax: +977-1-5553798

E-mail: info@npponepal.gov.np, Web : www.npponepal.gov.np

National Plant Protection Organization-Nepal

2082

GUIDELINES FOR SURVEILLANCE AND RISK ANALYSIS OF PLANT PESTS IN NEPAL

Endorsed by National Plant Protection Organization-Nepal, on 2071/01/22 BS,
First updated on 2081/02/29, Second updated on 2082/02/20



Government of Nepal

**Ministry of Agriculture and Livestock Development
Plant Quarantine and Pesticide Management Center**

Hariharbhawan, Lalitpur

Tel.: +977-1-5524352, 5553798 Fax: +977-1-5553798

E-mail: info@npponepal.gov.np, Web : www.npponepal.gov.np

National Plant Protection Organization-Nepal

2082

GUIDELINES FOR SURVEILLANCE AND RISK ANALYSIS OF PLANT PESTS IN NEPAL

Publication Year : 2081/82

Publication copies : 500

Published by : Plant Quarantine and Pesticide Management Centre
Hariharbhawan, Lalitpur

Printed at : Ganga printers Ph No. 9851016337, 014569455

FOREWORD



Plant health, now-a-days, has become a major trade policy issue; knowledge of the health status of a country's agro-forest commodities has important applications. These include the development of robust quarantine policies and the management of endemic pests. Plant health problems affect society in many ways. Many areas of society may be affected by incursion of new pests, diseases and weeds into a community. The favorable health status of agro-forest industries provides a competitive advantage in accessing foreign markets.

Accession into WTO is taken as the era of export opportunities in Nepal. But the exports from Nepal have not expanded to the same extent as trade between the developed members. The developed countries have expanded exports by using the rules of the SPS Agreement. At the same time, the government of Nepal is experiencing pressure from her agro-forest producers and traders to use the SPS rules to exclude commodities that are posing threats to agro-forest industries within the country.

Nepal is still unable to provide an adequate description of the health (pest) status of her plant based industries, so we are in a disadvantage when negotiating access to foreign markets. Prospective importers of our agro-forest commodities will assess risk based on our knowledge of the pests' status within our country while we are seeking to export. Our trade partners interested to import our agro-forest commodities will assess the likelihood of introducing pests found in our country with the imported commodity. Prospective importers also assess our phytosanitary measures being practiced to reduce risk to an acceptable level. Extensive specimen-based records are the key for us to negotiate with developed countries on a fair trading system.

If we want to expand exports of agro-forest commodities under the rules of the WTO, we must have the wide collection of our pests in our reference laboratories like that of NARC, CAL, NAST and NHM. The development of specimen-based pest lists can be accelerated through structured surveillance programs, focusing on the pests that might be carried on the commodity to be exported. Pest collections are significant because they provide the most reliable evidence of the plant health status of a country. These records are the foundation for developing robust policies for domestic and international quarantine and for developing pest-management strategies at the field level. The collections have taken particular significance since the accession of Nepal into the World Trade Organization

(WTO) in 2004. This guideline has been prepared by the NPPO of Nepal with a view to helping plant-health professionals. This guideline is prepared for the scientists, working in the NARC or NAST; plant protection officers, working under the ministry of agriculture development; professors teaching in the universities or the foresters working under the ministry of forestry. This guideline will help them to undertake Pest Risk Analysis and devise survey program on pest detection and to transmit specimens to the laboratory for identification and preservation.

Best Regards



Bhoj Raj Sapkota

Chief,

Plant Quarantine and Pesticide Management Center and
Head, National Plant Protection Organization (NPPO) Nepal

ACRONYMS

ALPP: Area of low pest prevalence
CDO: Chief District Officer
DNA: Deoxyribonucleic Acid
DOA: Department of Agriculture
DPR: Department of Plant Resources
FAO: Food and Agriculture Organization
FNCCI: Federation of Nepalese Chamber of Commerce and Industry
AEC: Agriculture Enterprise Centre
GIZ: German Technical Cooperation
GPS: Geographic Positioning System
IPPC: International Plant Protection Convention
ISPM: International Standard for Phytosanitary Measures
MOALD: Ministry of Agriculture and Livestock Development
MOFSC: Ministry of Forests and Soil Conservation
NARC: Nepal Agricultural Research Council
NAST: Nepal Academy of Science and Technology
NEHHPA: Nepal Herbs and Herbal Products Association
NHM: Natural History Museum
NPPO: National Plant Protection Organization
PQPMC: Plant Quarantine and Pesticide Management Center
PFA: Pest Free Area
PRA: Pest Risk Assessment
SPS: Sanitary and Phytosanitary Measures
RPPO: Regional Plant Protection Organization
VRM: Village Rural Municipality
WTO: World Trade Organization

Table of Contents

Part -1

Background	1
Terminologies	2
Designing a detection survey	4
A. Choosing a title	4
B. Reasoning for survey	5
C. Find the investors for survey	5
D. Identify the target host(s)	5
E. Identify the pests of the targeted host	7
F. Review the earlier survey plans	9
G. Select the survey site	9
H. Identify time for survey	13
I. Plan for required data	14
J. Identify the collection procedure	16
K. Label the specimens	18
L. Transport the specimens	19
M. Identify the human resource	20
N. Take required equipments to the field	21
O. Perform the survey	23
P. Analyse data	23
Q. Report the results	23

अनुसूची-१

PEST INFORMATION COLLECTION FORMAT FOR NATIONAL PEST DATABASE OF NEPAL	27
--	----

अनुसूची-२

FORMAT FOR SPECIMEN FORWARDAL	31
-------------------------------	----

अनुसूची-३

PEST INFORMATION REPORTING FORMAT FOR NATIONAL PEST DATABASE OF NEPAL	33
---	----

Part -2

WORK FLOW FORMATS FOR PRA

फाराम नं.-१

INFORMATION REQUIRED FOR PEST RISK ANALYSIS FOR IMPORT OF PLANT AND PLANT	37
---	----

MATERIALS INTO NEPAL

	फाराम नं. २	
INFORMATION NEEDED TO UNDERTAKE PRA IN NEPAL		39
	फाराम नं. ३	
FORMAT FOR PEST RISK ANALYSIS		41
FULL PEST RISK ANALYSIS		45
	फाराम नं. ४	
ALOP FORM		62
	फाराम नं. ५	
FORMAT FOR PRODUCT/COMMODITY RISK ASSESSMENT		63
	फाराम नं. ६	
शत्रुजिव जोखिम विश्लेषणमा आधारित अन्तरराष्ट्रिय पहुंचको प्रस्तावका लागि निवेदन साथ संलग्न गर्नुपर्ने विवरण		67

PART -1

BACKGROUND

Accession to WTO is taken as the era of export opportunities in Nepal. But the exports from Nepal have not increased to the same extent as trade between the developed members. The developed countries have escalated exports by using the rules of the SPS Agreement. At the same time, the Government of Nepal is obliged to use the SPS rules to exclude commodities that are posing threats to the related industries within the country. Nepal should provide an adequate description of the health status of plant based industries, while negotiating access to foreign markets. Prospective importers of Nepalese forest related commodities assess risk of introducing new pest based on the authentic pest information provided. Prospective importers also assess the phytosanitary measures being practiced in Nepal to reduce risk to an acceptable level. Extensive specimen-based records are the key for Nepal to negotiate with importing countries on a fair trading system. This document gives detailed guidelines for detection surveys of the plant pests in natural agro-ecosystems, and is basis for specimen based records to be developed by the NPPO.

This document describes the components of survey for the purpose of pest detection in an area. The data generated from the survey is supplied for pest risk analysis. The data also supplies information to establish pest free area and to declare ALPP. To increase exports of forest related commodities under the rules of the WTO, there should be the wide collection of pests in the laboratories as of NARC, NAST and NHM. The development of specimen-based pest lists is made through well planned survey programs, focusing on the pests related to commodities in trade. Pest collections provide the most reliable evidence of the plant health status of the country. The collection and records of pests are the foundation for developing robust policies for domestic and international quarantine and for developing pest-management strategies at the field level. This guideline is prepared for the plant health scientists, plant protection officers, researchers and for the foresters working under different organizations. This document covers the procedures in detecting arthropod pests and plant pathogens and weed in plantations, forests and natural ecosystems. This document has included and followed the ISPMs whenever possible. This document is not intended to guide the technicians to detect the pests in the consignments at the entry or exit points.

Under Plant Protection Act 2064, Clause 6 (2), survey and surveillance function and responsibility is designated to NPPO as per the sub clause (i). “To perform such other function as prescribed”. Plant Quarantine and Pesticide Management center is authorized

NPPO. This technical guideline to undertake survey program related to pest detection has been prepared by the NPPO of Nepal. This document also guides to transmit specimens to the laboratory for identification and preservation. The publication covers the planning of survey programs for building specimen-based lists of pest. The guideline helps design a survey program, emphasizing the need to carefully document the process. The guidelines also provide advice on how to approach the critical issues of designing a statistically valid program that will meet the most rigorous demands of trading partners and others who must have faith in the results. The glossary of ISPM terms that relate to surveillance is published in ISPM No. 5. The most relevant entries are reproduced in the terminologies of this guideline.

TERMINOLOGIES

Survey: Survey is an official procedure conducted over a defined period of time to determine the characteristics of pest population occur in an area. It includes regular systematic collection and identification of major pests, diseases and weeds as well as their natural enemies.

Surveillance: Surveillance is an official process which collects and records data on pest occurrence or absence by detailed study of the pests and beneficial insects. In other words pest surveillance monitors closely and regularly the development of plant pests, their natural enemies and other prevailing factors in an ecosystem.

Specific surveys: Procedures by which NPPO obtain information on pests of concern on specific sites in an area over a defined period of time. Pest list survey is one of the types of specific surveys within detection survey.

Detection survey: Survey conducted in an area to determine if pests are present [FAO, 1990; revised FAO, 1995] Note: The purpose of detection survey is to detect the presence or absence of the pest in a given area or production sites. These are more frequently carried out to determine pest status in an area and they follow a definite survey plan, which is approved by the Plant Quarantine and Pesticide Management Center (PQPMC)/ NPPO or any other institute or organization authorized by PQPMC and are systematically organized. These surveys are carried out either seasonally or annually and/ or following the eradication measures applied to a pest in a given area or production sites. These surveys are organized following a definite survey methodologies based on statistical sampling, which are determined after taking into account the biology of the pest and employing appropriate detection techniques such as field diagnostic kits, traps etc. The results of survey are documented and communicated.

General surveillance: A process whereby information on particular pests which are of concern for an area is gathered from many sources, wherever it is available and provided for use by the NPPO.

International Standard for Phytosanitary Measures (ISPM): An international standard adopted by the Conference of FAO, the Interim Commission on Phytosanitary Measures or the Commission on Phytosanitary Measures, established under the IPPC.

National Plant Protection Organization (NPPO):

Official service established by a government to discharge the functions specified by the IPPC. The IPPC (1997), in relation to its main purpose of “*securing common and effective action to prevent the spread and introduction of pests of plants and plant products*, (Article I.1) requires countries to make provision, to the best of their ability, for an official national plant protection organization,” (Article IV.1) whose responsibilities include the following: “*...the surveillance of growing plants, including both areas under cultivation (inter alia fields, plantations, nurseries, gardens, greenhouses and laboratories) and wild flora, and of plants and plant products in storage or in transportation, particularly with the object of reporting the occurrence, outbreak and spread of pests, and of controlling those pests, including the reporting referred to under Article VIII paragraph 1(a)...*” (Article IV.2b). **ISPM 17.**

Pest: Any species, strain or biotype of plant, animal or pathogenic agent injurious to plants or plant products.

Pest free area (PFA): An area in which a specific pest does not occur as demonstrated by scientific evidence and in which, where appropriate, this condition is being officially maintained.

Pest record: A document providing information concerning the presence or absence of a specific pest at a particular location at a certain time, within an area (usually a country) under described circumstances.

Pest risk analysis (PRA): The process of evaluating biological or other scientific and economic evidence to determine whether a pest should be regulated and the strength of any phytosanitary measures to be taken against it.

Pest status (in an area): Presence or absence, at the present time, of a pest in an area, including, where appropriate, its distribution, as officially determined using expert judgment on the basis of current and historical pest records and other information.

Blitz surveys: The purpose of blitz surveys is to detect all pests present, even those in low numbers, and to identify less visible symptoms and newly emerging pests. These surveys involve the intensive inspection of all plants in a given field site or at a set time, generating pest lists for a host or range of hosts.

DESIGNING A DETECTION SURVEY

The collection and recording of pest information is fundamental to all WTO/SPS and trade related activities. NPPO should be in a position to validate declarations of the absence or limited distribution of pests. Particularly, NPPO should decide the objective and procedure of any specific survey to meet the country's requirement. It should guide the surveyors while collecting information about a pest in the field. It should provide the overall framework for planning any type of specific survey. NPPO should also develop specific protocols for survey of specific pest. Specific surveys an official survey and so should follow a plan which is approved by the NPPO. Depending upon the bilateral trade requirements, both exporting and importing country's NPPO are in general, responsible for determining the process and the techniques of gathering the information on pest. When planning a new survey, the details of the design need to be carefully recorded and justified. Reasons and decisions need to be justified as the result of the survey requires approval from the NPPO. The survey plan should include:

- Properly defined purpose of survey (early detection, assurance for pest free areas, development of pest list, decision to apply phytosanitary measures, finding the extent of pest spread, biosecurity and incursion management, etc.)
- Identification of the target pests (their life cycles and identifiable characteristics, pest vector, host, etc.)
- Identification of scope (such as geographical area, production system, season)
- Identification of timing (dates, frequency, duration)
- In the case of commodity pest lists, the target commodity
Indication of the statistical basis (such as level of confidence, number of samples, selection and number of sites, frequency of sampling, assumptions).
Description of survey methodology and quality management including an explanation of sampling procedures, sample collection and laboratory analysis, diagnostic procedures, and reporting procedures.

A. CHOOSING A TITLE

- The title should be simple, short and imparting the subject of survey.
- Name, address and the contact number of the person responsible for producing the survey plan should be clearly written, so that the confusion happening in the field can be resolved by the survey team after contacting the planner.

- Name, designation and affiliated institution of survey team leader and composition of survey team
- Name and addresses of the institution and collaborating partner/s
- Defines period of time (days, weeks, months or years)
- Command potential survey areas or locations
- Estimated cost for survey

B. REASONING FOR SURVEY

There might be several reasons for pest survey. The survey plan should clearly explain the reasons for pest survey. The reason may include development of specimen based pest list and host list, to obtain data to support pest free areas (PFAs), pest free places of production (PFPP) or pest free production sites (PFPS), determination of pest status in an area, guidelines for pest eradication program, pest reporting, requirements for the establishment of areas of low pest prevalence, etc.

C. FIND THE INVESTORS FOR SURVEY

Survey needs adequate funding and therefore necessary to approach suitable investors/donors...Preparation of the detail budget plan for survey is essential.

D. IDENTIFY THE TARGET HOST(S)

The importance of the host for nutritional or medicinal value and its significance to small communities, and its national or regional economic importance should justify the effort of the survey. The growth habits of host and any aspects of life cycle that are relevant to the diagnosis of the pests to be investigated should be recorded. Growing condition, growth habit, accessibilities to different parts, location characters, host's distribution pattern, and regional distribution, etc. should be clearly described before undertaking surveys. In applying for the PFA, the survey should provide information on the location and pattern of host plants distribution within the area of interest.

Host names:

List the common and scientific names of targeted host plants.

Value of host or commodity:

Describe the importance of the hosts; for example, their nutritional or medicinal value to local communities, and their national or regional economic importance.

Growth habits and life cycle of host plants:

Describe the growth habits of each host and any aspects of their life cycle that are relevant to the diagnosis of the pests to be investigated. List how and at what condition the host plants of interest are grown. Describe how tall and bushy does the vegetation grow, how much of the plant could be seen and accessed, ability to collect a specimen from the crown, the middle near the main stem, the tips of the growth, or at the base of the plant, etc.

Accessibility of the host plants:

While designing a survey, consider the vegetation and the areas in which the pests are to be surveyed. Information about the accessibility of hosts is important for the future surveyor using the report as part of general surveillance; it should explain to the future surveyors why only certain places were surveyed. The plan should explain in detail on how the host plants are ordered? And ability of surveyor to walk through the row or between the plants, could the surveyor see the entire plants in a row or in random? Where can the surveyor walk? How much damage caused by walking through the crop would be accepted by the property managers? How far the surveyor can expect that someone could see into the crop or forest? , what is the terrain like? Are there remote parts? Are there any dams, rivers or fences that may affect the accessibility of the site? etc.

Regional distribution of the host:

A list of properties and sites with host plants should be developed from a number of sources, including industry and government records and personnel, local grower groups and cooperatives, packing operators and distributors, extension staff, researchers and property owners. Aerial photographs can be useful for identifying areas that are densely populated by hosts, such as production areas. The survey plan should consider all alternative hosts, as well as the susceptibility of endemic flora in forest, parklands, gardens and other areas close to the detection site.

Describe the distribution of the host in the country/region of interest. List all of the locations by name. Before going for the pest status survey, it is essential to have a report of host distribution survey performed earlier. If possible, consider the host distribution report of the departments under the Ministry of Forest. In applying for the PFA, the survey should provide information on the location and pattern of host plants distribution within the area of interest.

E. IDENTIFY THE PESTS OF THE TARGETED HOST

In case of pest-host specific survey, i.e. finding out the status of specific pest in specific host; or in case of known pests (more than one pests) are to be surveyed in a specific host, the following information on the specific pests of targeted host are required in the survey plan:

Find the sources of information:

Find the information on pests about their life cycles and identifiable characteristics by reference consultations or by browsing on the credible websites. The information sources may include: NPPOs, other national and local government agencies, research institutions, universities, scientific societies, producers, consultants, museums, the general public, scientific and trade journals, unpublished data and contemporary observations. In addition, the NPPO may obtain information from international sources such as, IUCN, Regional Plant Protection Organization (RPPOs), etc. Other sources may include: existing PRA reports; either conducted by NPPO of Nepal or by agencies of other countries, reference collections of insects/ pests and plant pathogens of economic importance, and the pest interception databases from quarantine authorities. ISPM 8 has a basis for evaluating the reliability of a pest record that could equally be applied to assessing information sources to be used in developing the survey plan.

Record the Pest names:

Begin by creating a list of the scientific and common names of the pests that attack the host together with synonyms wherever applicable.

Record the Pest vectors:

Identify any vectors of the pests to be surveyed. If the pests have vectors, include them in the list of target organisms.

Record the possible Pest impacts:

Consider why these pests are chosen, are they regarded as major pests or pest threats? Do trade partners want more information on the status of specific pests in the targeted area? In general terms, describe how the pests would be likely to affect a host, production system or ecosystem, and the industry as a whole.

Pest characteristics:

The diagnostic characteristics of a pest, or symptoms of its presence, should be compiled. It is necessary to create a team of specialists and establish laboratories that have experience with the pests and the diagnostic capacity to identify them, depending on pests intended

for survey. Wherever host plants are involved, describe the parts of the plants most likely to be infested or infected, and parts of the plant that should be examined. Does the pest target a commodity; such as fruit or modified root? Is the pest associated with particular stages of a host plant's growth? Is the pest attracted by light or pheromones? Describe where the pest or the characteristic symptoms would be found on the host or commodity; for example, flying above a plant, bored into bark, the underside of leaves, at the base of plant, presence of curly leaves, etc. Are there any factors that affect symptom development, such as sub-species, growth stage, season, pesticide application and climatic conditions? Identify all the likely sites that the pest might have infested. Research the epidemiology of the pest, its means of survival, reproductive rate and life span. Include all available information about the pest's life cycle. In summary, prepare pest information sheet. Pest information sheets provide identifying details of pests that the survey team can refer to during the field survey.

When applying for PFPP and PFPS, the survey should provide the following essential information:

Natural spread habit: fast or slow, spreading long or over short distances;

Artificial spread habit: limited or high chance of the pest being spread artificially; Host range: wide or limited;

Survival rate between seasons: high or low;

Rate of reproduction: slow or moderate or fast;

Detection: easily or hardly detectable;

Control measures: effective, practical and easily available or no measures.

Collecting reference specimens and images:

Having handout material that can be used in the field may be critically important, particularly if the pest has not been seen before by the surveillance team. Having a reference collection of pressed samples of plants or affected plants, or small specimen collections of invertebrate animals may also prove useful as long as they are not cumbersome and can be protected from damage. Electronic images or photographs of the concerned pests can be used as reference images. These can be used to create pest information sheets.

Alternative hosts:

Finding out the distribution of known alternate/collateral hosts of a particular pest in an area is called pest specific host distribution survey. It is done for delimiting the area of infestation by a particular pest. Listing of entire host range of a particular pest, including vectors, and surveying for the presence of hosts in an area is particularly important for declaring and managing PFA or PFPS or PFPP or for deciding any form of incursion management or for determining the buffer area where internal quarantine system would

be implemented. Alternate/collateral host survey of specific pest is done with the objective to locate and destroy the hosts before declaring the PFA/PFPS or PFPP.

F. REVIEW THE EARLIER SURVEY PLANS

NPPO of Nepal and the concerned officials in NPPO can have existing plans or can provide the contact with others who have designed survey plans. If the plan is connected with trade, the NPPO needs to involve as part of the process.

G. SELECT THE SURVEY SITE

There are usually six steps involved in site selection:

- Select the ‘Province’.
- Select the ‘district(s)’ involved
- Select the ‘local level’ in the districts that could be surveyed.
- Select the ‘field sites’ within each place: forests, host communities, ports or markets.
- Select the ‘sampling sites’ within each field site. This could mean, individual plants, trees or produce, transects, or trees to which pheromone traps could be attached.
- Select the ‘sampling point’, which is relevant to choose specimens within a sampling site.
- For undertaking detection survey for PFPP and PFPS, the following acceptable characteristics of the site need to be fulfilled:
 - it should be a single property
 - it should have clearly defined boundaries, including any buffer zones
 - it should be sufficiently isolated from possible pest infestations
 - there should be no other known hosts within the boundaries, including the buffer zone.

Methods of site selection:

There is no single best method for site selection. The main point is to transparently document the choices and reasons for such choice. The method can then be considered and discussed with other parties for agreement with the basis of the choice, given the circumstances.

Select the appropriate sites for survey:

Full sampling and blitz surveys are always recommended and universally accepted methods of providing the most detailed data for detection surveys in random population. If it is not possible to cover all sites at each level, it is necessary to select the site to cover. . To do this, use one or a combination of the following four tools:

1. Random sampling: This involves assigning all sites (of the uniform level), a number or symbol and then by using a random number generation method, the sites are selected and recorded.
2. Systematic sampling: This involves selecting criteria to divide the sites into some form of regular intervals and then selection of sites accordingly. . For example, surveying every second site when listed by name in alphabetical order, setting up a grid of traps or parallel transects of a site.
3. Stratification sampling: This method can be used in combination with random sampling. This involves dividing the sites into logical categories and then systematically or randomly choosing sites from within the categories. Example: 20 villages (level: place) are to be surveyed for diseases of *Sapindus mukorossi* (Ritha). Each village has 15 production sites (level: field sites), a total of 300 production sites. If 100 farms are to be surveyed, randomly choose the 100 from all 300. By chance, this may result in some villages having all their production sites surveyed and others having none. If it is important that all villages be surveyed, the selection of the 100 sites can be stratified by village such that, for example, five farms per village are chosen randomly.
4. Targeted site selection: The sites are chosen based on where the pest is most likely to be, thereby deliberately biasing the selection process in favor of finding the pest. Targeted sites can be in the field or forest where surveillance is focused on host plants or sites where the pest is most likely to be present. Field workers, property managers and others working at the places of interest may be able to provide local knowledge of where pests present may have been observed. With this method, it is possible to identify particular niches where the pests could be found.

Random number generation

Randomizing the order in which sites are visited, can be achieved by assigning each site a sequential number and carefully listing the sites and their numbers. Cards labeled with site numbers or names can be well shuffled and read off in the order they appear. These methods are useful unless the number of sites is more than a literal handful of cards. Using random number tables is the best of all methods of randomization. It is important to record the order of the sampling sites as species-accumulation curves need to be plotted to determine whether the survey is completed for detection surveys.

Species accumulation curves for detection surveys


After looking at a number of quadrates, the number of new species added to the list will be fewer and the increasingly smaller amounts of information gained need to be weighed against the effort. The sequential sites should be chosen by random selection so that they are unlikely to cluster together. Then the field sites should be stratified into sections.

A species accumulation curve is used to determine the number of sampling sites needed to survey. The process requires recording the number of new pests collected at each new site, then plotting the accumulated number of pest species—with sites across the X- axis and number of pests along the Y- axis. The number of new species will eventually decline with the increasing number of sites examined. A curve is then plotted to the data points. When the curve has flattened for example, five consecutive sampling sites, if no or few species are added with each additional site, the survey is complete.

This exercise should be repeated in different production areas or districts if there is reason, such as climatic variations, to suspect that the pest list may vary from site to site. Species accumulation curves can also be drawn for one location but over time. This means that surveyor would plot the number of new pests on the Y- axis against time intervals on the X- axis. Surveyor may wish to do this, if the pest distribution on a given host is seasonal.

Plan to install the static traps for insects for detection and monitoring

Insects can be caught by static traps that attract insects by light, color or pheromones. The insects are then removed from the trap and identified. These traps are useful primarily for identifying whether or not a pest is present in the area. The sitting and density of traps is critical. Sitting and density are determined by the trap type and the manufacturer's instructions, and applied according to the survey setting. Traps are often used to estimate the prevalence of pests in the area. In some cases, the number of insects trapped is directly proportional to the true pest prevalence.



(Source: Mc Maugh, 2005)

Fig. 1: Species accumulation curve for detection survey

Calculating sample size

Some main statistical parameters need to be understood by the planner. Parameters like design prevalence or confidence etc. is expressed in percentages except for sample size which is in whole numbers.

Design prevalence

This is estimated prevalence (detection threshold) and usually based on a pre-survey estimate of the actual prevalence of the pest in the field, and used to determine the sample size. Clearly, for area freedom, the design prevalence and actual prevalence of a pest are expected to be near-zero. For surveys that monitor a pest that is known to be present, the design prevalence can range from near-zero to 100%. If the design prevalence greatly over-estimates the actual prevalence, the sample size calculated will be too small to detect the actual prevalence. If the design prevalence under-estimates the actual prevalence, then the sample size will be larger than needed to detect the actual presence, leading to over-sampling. If NPPO is unable to predict a meaningful prevalence, especially in case of detection survey to be done for the first time, it is necessary to choose a prevalence level that is acceptable to all parties. For detection survey, the design prevalence is usually taken as 1%.

Confidence

Statistical confidence is the probability that the actual prevalence will be within the range of the design prevalence. The relationship between confidence and sample size is simple; the more sites surveyed the more certainty of the accuracy of the estimated prevalence. As a general rule, a detection threshold of at least 95% confidence is considered acceptable. Confidence up to 99.9% can be necessary in some instances, for example in declaring PFA. Trading partners may require a particular level of confidence that the pest would be detected in a survey, independent of any logistical or financial constraints.

Sample size for detection surveys

Sample size is the number of sites that need to survey in order to detect a specified proportion of pest infestation with a specific level of confidence, at the design prevalence. Formulas for calculating sample size are given below. These formulas are used when the survey is designed to detect a pest, and where the actual prevalence is likely to be rare.

A simple relationship exists between sample size, confidence level and detection threshold, where confidence is expressed as a percentage and detection threshold on a scale between 0 and 1.

Formula: Confidence level = $1 - (1 - \text{design prevalence})^{\text{sample size}}$

And therefore,

$$\text{Sample size} = \frac{\log(1 - \text{confidence level})}{\log(1 - \text{design prevalence})}$$

The following table presents the calculations performed on the basis of these formulas:

Confidence	1 – confidence	Design prevalence	1 – design prevalence	Sample size
0.95	0.05	0.01	0.99	298
0.95	0.05	0.02	0.98	148
0.99	0.01	0.01	0.99	458
0.99	0.01	0.02	0.98	228
0.95	0.05	0.001	0.999	2,994
0.95	0.05	0.002	0.998	1,496
0.99	0.01	0.001	0.999	4,603
0.99	0.01	0.002	0.998	2,300

(Source: Mc Maugh, 2005)

Accuracy of methods

This deals with how well the survey will detect a pest when it is present. For example, if it is required to look for the pest on a row of trees but due to some reason the surveyors plan to inspect only along a straight line then the method of inspection is not 100% accurate. It is expected that the observer cannot see all of the trees, if the foliage is dense or the symptoms or pests are not obvious. In such case, 80% can be specified as the accuracy of this method. Some methods can reasonably be expected to be 100% accurate. So if the surveyors are not going to follow the exact protocols of the observation, because of several reasons, then the sample size should be readjusted.

Adjusted sample size = (sample size above)/method accuracy size (generally 0.8).

H. IDENTIFY TIME FOR SURVEY

Pest survey:

The survey should be performed when the pest is most likely to be present and in an identifiable state. The timing of survey procedures is determined by: the life cycle of the pest, the phenology of the pest and its hosts, the timing of pest management programs, whether the pest is best detected on plants in active growth or in the harvested state (ISPM 6). If the best time to survey is not known for the pests, start by finding out its seasonal habits. At what stage of the host's life cycle do the pests prefer to infest it? How long will it persist? Does it die back; such as during the wet or dry season? Can it survive if the host is dead or dying? Does normal management of the forest control or eradicate the pest? How quickly does it reproduce and spread? How long does the pest survive under

different conditions and at different stages of its life cycle? Are there weather conditions or weather events that may influence the pest's life cycle or survival? Other factors that may determine the timing of survey are:

- when the pest is most active
- accessibility and availability of vehicles
- time of local festivals or community events
- time of sowing, seedling emergence, flowering, fruit maturation and harvesting of hosts
- time of flowering for weeds
- time of obvious symptoms

Host survey:

The timing of a survey is particularly important when developing pest lists, as it is critical that host plants are examined throughout their life cycle since different pests prefer different stages of the host development. The minimum stages of development that should be surveyed are:

- seedling emergence
- vegetative flushing stage
- flowering stage
- fruiting stage
- post-harvest stage
- Consider examining the host plants under different weather conditions.

Frequency of the survey

Some surveys need to be performed several times. For example, every 2 weeks when managing a pest in a plant species, or annually during harvest to support a pest-free-area status, or according to periods in the pest's life cycle. If trading partners are involved, the frequency of survey is based on mutual agreement. . Also, it is essential to revise the timing and frequency if they are dependent on weather conditions or events. The frequency of survey also depends on the trading partner's need. The importing country may require that pest free area status be verified for 'one or more years' before the year in which export would commence, or simply from the year of export onwards. The frequency can be adjusted according to the perceived risk of the site.

I. PLAN FOR REQUIRED DATA

Tagging the site:

It is necessary to mark sampling sites in the field, even if, it is not intended to return to the same site. It is possible that a specimen or observation taken could be lost or destroyed.

So with careful notebook entries and a marked site, the surveyor would be able to revisit the site, if needed. The quality of tags and ink should be able to endure a variety of weather conditions.

Recording site details:

The location and unique identifying details of each site should be recorded in a notebook. Describing the sampling site would include information such as a GPS reading, a unique number, distances from visual cues, number or nearest number of plant in a row, or any distinguishing topographical features.

Field data to be recorded:

The most important is to record any information that could otherwise be forgotten, such as the dates of survey, the weather at the time, the site details, the names and contact details of the local informants involved, variations in the team member involved in survey and any other relevant details. While recording the data one should fill up the form provided in the Annex “1” that allows for recording all the information required for NPPO. For PFPP and PFPS detection survey ‘may be required’ of the harvested commodity at the production site. In this case, the additional information required depends on the demand of the importing country.

Units for data:

Data are normally reported in terms of a unit of measure, usually the number of pests per unit area. The number might be a direct count of the pests or could be a scale of intensity of the pest that is recorded. For early detection or to support pest-free-area status, the pests will rarely be found. The pest count usually is zero, but it is still important to quantify the amount of effort expended for statistical purposes. For example, ‘600 trees were examined in each of 20 farms in an area, with no evidence of the pest’. In cases where the pest is numerous, or particularly for symptoms of plant pathogens, whole numbers of pests are not possible or useful. Instead, a scale of cover of the host or a standardized measure of the pest could be used.

Proving the negative data:

It is very important to record negative data, i.e. locations surveyed where the pest was not observed, so that there is a record of the effort expended to look for the pest. It is particularly important in surveys to support pest-free-area status. The validity of negative records depends on a number of factors:

- the pest is known to produce easily noticed signs or symptoms
- the host species is widely distributed and has high population levels
- the host is economically important and is likely to have been examined by

plant protection specialists

- the pest is relatively easy to identify
- Environmental conditions are conducive to infection and pest development.

J. IDENTIFY THE COLLECTION PROCEDURE

Pest specimens should be collected and handled with the best possible care to preserve the diagnostic features for identification. If specimens are to be sent away for identification, consider collecting two or more specimens and preserve them adequately. One is to keep with surveyor and one is to send for identification. After identification, surveyor should provide the specimen to NPPO and that will be used for future reference. Specific protocols developed by NPPO for specific pest should be used for collection, handling and preservation. Insect and mite pests may be collected by netting, beating, aspirating and vacuuming, trapping, extracting, specialized collecting, etc. Similarly, there are several methods of getting disease specimen from different parts of the plant.

Basic specimen collection procedures:

- Sterilize any implements with 70% ethanol or 0.5% available chlorine solution before and after each sampling.
- If considered to be a root problem, include soil and crown (lower stem) tissues with root samples.
- It is essential that the time between sampling and dispatch of the sample for identification be kept to a minimum.
- If possible, sample from perceived area of minimal damage to perceived high damage within field and on individual plant

Insect samples (use specific protocols provided by NPPO):

- Where possible, collect a large number of specimens of all life stages.
- Collect specimens with appendages such as antennae, wings and legs.
- Use leak-proof alcohol and resistant plastic container with screw-top lid.
- If sending small and/or soft bodied insects (example: thrips, aphids, mites and larvae), place specimen in 65% ethyl alcohol and completely fill the container
- Do not remove mealy bugs or scale insects from the leaves or stems on which they are feeding as this will damage their mouth parts and make identification difficult. Instead, cut out leaf tissue around the insect and place this in alcohol.
- While sending hard-bodied insects (example: beetles, moths, grasshoppers and fruit flies), carefully fold specimen in tissue paper and place in crush-proof plastic tube or container with several holes in the lid for ventilation.
- Retain and store a spare sample in a secure, cool and dark location.
- If possible, store sample in freezer for 2 hours before dispatch to kill the insect.

- Clearly label all samples.
- Do not send live insects unless live sample is demanded by diagnostic laboratory.

Fungus and Bacteria (use specific protocols as provided by NPPO):

- Try to select the sample on the same day it is to be sent, to ensure freshness.
- For fungal and bacterial samples, store under 2–5°C until it is sent, but note that some pathogens do not survive in cold conditions.
- Select samples at the margin between the diseased portion of the plant and the healthy portion.
- Select a fresh, representative and generous sample covering the full range of symptoms.
- If considered to be a root problem include soil and crown (lower stem) tissues with root samples.
- Place samples in self-sealing ventilated plastic bags with some dry tissues or paper towel to absorb excess moisture.
- While submitting a succulent sample, wrap in dry tissues or paper towel and pack firmly in a crush-proof container.
- Do not send dead plant material.

Nematodes:

- Sampling of soils that are very wet or very dry should be avoided.
- The soil for the sample should be taken at least 5–10cm below the surface.
- Separate samples should be taken from the badly affected and normal areas
- Trees and vines should be sampled at the drip circle.
- Individual sample size should be about 250–300g.
- Roots should be either included in the sample or taken separately – about 25–100g, taken at random.
- Affected stems or leaf material should be kept separate from soil and/or root samples.
- Samples should be kept cool – do not leave in the sun or in a closed vehicle left in the sun.
- If immediate dispatch or processing is impossible then samples can be stored in a refrigerator at 4–8°C for several days.

Phytoplasmas and Viruses:

- Plant material that is suspected of being infected with a virus should be collected, cut them into pieces and place in a plastic container containing calcium chloride (CaCl₂) crystals or silica gel, but separated by cotton wool.

- Sterilize the scissors or safety blades in alcohol or a 10% sodium hypochlorite (NaOCl) solution between samples to prevent cross contamination.

Weeds:

- Select vigorous specimen, avoid insect or disease-damaged plants.
- Collect sufficient material to fill a herbarium sheet (ca. 450 × 300 mm) and still leave enough room for the label. Plants too large for a single sheet should be divided and pressed in a series of sheets.
- Where possible, include mature leaves, juvenile leaves, buds, flowers, fruits, and bark.
- Use specific protocols for collecting and pressing for trees, bulbs, rhizomes, cones, palms, vines, grasses, rosette plants etc.

Pressing and care of specimens:

Specimens should be pressed as quickly as possible after collection. If this is not possible, specimens can be stored in plastic bags, preferably wrapped in damp (but not wet) papers. Bags should not be packed tightly, and should be kept cool and moist. Make sure that each bag is correctly labeled. Place each specimen, with numbered tie-on tag attached, in a fold of several sheets of newspaper, and place in the press. While filling the press, try to keep it level to allow even distribution of pressure. Close the press and exert pressure with the straps. The plants in the press should be dried fairly quickly, in a warm place. The specimens must not be left in damp papers otherwise they will go mouldy. It is therefore necessary to go through the press daily during the first few days and change the plants into dry newspapers. Then continue to inspect press daily and change newspapers as necessary until the plants are dry. Delicate plants and petals may be lost in changing and should be kept in tissue-paper.

K. LABEL THE SPECIMENS

Label the specimens in the field, at least in a temporary way until a full and appropriate label can be made. The following information should be included in the label:

- scientific name of host and plant parts affected;
- scientific name and life stage or state of pest;
- family/order of the pest;
- locality, such as location codes, addresses, coordinates;
- Date of collection and name of collector.

To submit specimens to a diagnostic laboratory or expert for identification, ask them about the required condition of specimen and format of details that must be submitted together. Annex 2 provides the general format to fill up before sending the specimen to diagnostic laboratory. Specimen details should be written with water/alcohol proof inks. If the specimen is in a jar or container, the jar itself should be best labeled rather than the

lid. Microscope slides should be labeled with small stickers on the upper side of the slide, away from the specimen itself.

Labelling of pest specimens

PINNING & LABELLING INSECT WITH COLLECTION DATA

ALWAYS label specimens with complete collection data in or on each container. Label should contained collection data information in a label

• SCIENTIFIC NAME	<u>Location label</u>	<u>Host plant label</u>
• COMMON ENGLISH NAME	Country: District	ME trap
• HOST OF INSECT	Date: 4.xi.2022 (ex.)	Citrus orchard
• PLACE	Collector's name	Reared from
• DATE OF COLLECTION	<u>Identity label</u>	<i>Citrus sinensis</i>
• NAME OF THE COLLECTOR	Scientific name of Insect	
• ALTITUDE OF THE PLACE	Ex. <i>Bactrocera dorsalis</i> (Hendel)	
• GPS POINT	Det. Author's name	
	Ex. D. Adhikari	
	Month, Year	
	Ex. Dec. 2022	

Labelling of insect specimens

Location label	Host plant label	Identity label
NEPAL: Sindhuli 4.xi.2022 D. Adhikari	ME Trap Citrus orchard	♂ <i>Bactrocera dorsalis</i> (Hendel) det. David, K. J., Dec., 2022
	CL Trap Citrus orchard	♀ <i>Zrugodacus cucurbitae</i> (Coquillett) det. David, K. J., Dec., 2022
	Reared from fruit of <i>Citrus sinensis</i>	♂ <i>Bactrocera minax</i> (Enderlein) det. David, K. J., Dec., 2022

Note the way the dates are written:
day, month, year
with month always
in lower case
roman letter

Collection by
sweeping on a host
plant,
Or
Found resting on a
host plant

L. TRANSPORT THE SPECIMENS

If the surveyor him/herself is transporting the specimens, it is necessary to ensure that they are being properly protected. If the specimens need to be sent by postal service, greater care should be taken in packaging them to cope with possible mishandling during transport. Take extra care when dealing with the following:

- **Live pests:** These require ventilation, so ensure that air can get in and the pest cannot get out. Keep plant specimens alive by wrapping in slightly damp paper and sealing in a plastic bag. Ensure that the specimens are protected from extremes of

temperature during the journey.

- **Glass or breakable containers:** These should be packed carefully so that the glasses do not touch each other or hard surfaces and break. Such containers can be protected by packing them into a second container that is at least 2.5 cm larger on all sides, with packing material placed in the gap.
 - **Multiple specimens:** If two or more specimens are to be packaged together, make sure that each is well labeled.
 - **Specimens preserved in alcohol:** The containers need to be leak-proof.
 - **Timing:** Submit specimens as soon as possible after collection.
 - **Postal or courier service requirements:** Check whether the postal or courier system has restrictions on sending particular volumes of alcohol, pests, container types or anything relevant to the specimen to be sent.

M. IDENTIFY THE HUMAN RESOURCE

Personnel involved in surveys should be adequately trained, and where appropriate audited, in sampling methods, preservation and transportation of samples for identification and record keeping associated with samples (ISPM 6). It is essential to think about the formation of the survey team involved in the field. Consider how experienced they are in recognizing the pest and if they need additional trainings. The team should be informed about the whole process, including the standard methods to be adopted to identify and record pests. Consistency in diagnostic skills of survey team members should be conformed before going into the field.

NPPO is responsible for all type of surveys, inspections and any other systems needed to verify pest status. The surveys should be carried out by NPPO personnel, or by experts authorized by the NPPO. For PFPS, PFPP and PFA, NPPO must certify the management, technical and operational skills of the producer to prevent the pest entering the place or site and their ability to manage the pest if it was detected on site. NPPO should provide the producer with trainings in pest-management systems when necessary. NPPO is also responsible for checking the regulations of the importing country and assisting the producer in establishing conditions that would lead to compliance with such regulations.

N. TAKE REQUIRED EQUIPMENTS TO THE FIELD

Below is the list of equipment to consider taking during field surveys. Depending on the objective and duration, the list of equipment can be reduced or increased.

Personal items:

- Light raincoat (in rainy seasons);
- Snake-proof boots, pants, and gloves;
- Drinking water and food; such as glucose in the form of hard candies etc.;

- Mosquito repellent;
- Watch;
- First-aid kit: standard items plus bite cream, para-cetamol, anti-diarrheal medication, anti-allergy tablets, antiseptic (like Dettol), rehydrating drink sachets (like Tang), chlorine water-purifying tablets (like Piyush); Mobile phone with SIM card;
- ID card.

Pest information sheets:

- Field data recording formats;
- Field guide.

Recording data:

- Water-proof/alcohol-proof pens;
- Field notebook;
- Water-proof paper may be needed to write on during rainy season.

Specimen-collecting equipment:

- Collectors tags;
- Plastic and paper bags;
- A magnifying glass/hand lens;
- Specimen tubes;
- Preserving alcohol (70–90% ethanol);
- Tissue paper;
- Para film;
- Tweezers/forceps/scalpel;
- Camera;
- Small pair of binoculars;
- Secateurs;
- Spade;
- Hand-held geographical positioning system (GPS) unit;
- Maps of the area;
- Compass;
- Diagnostic keys (identification, surveying, disease/pest rating scales);
- Permits for collection, survey and specimen transport;
- Pen-knife;
- Gardening gloves;
- Random number table or pack of cards, dice etc.;
- Calculator, stapler, staples;
- Lighter/matchbox;
- Disinfectant wipes;

- Measuring tape;
- Spray paint;
- Brightly colored ribbons/tape;
- Trowel or spade;
- Plastic zip-lock bags of various sizes;
- Axe;
- Portable icebox;
- Envelopes (Nepali paper- various size);
- Small buckets (to carry intact soil samples with plants);
- Survey bag-waterproof and non-leather;
- Hunting jacket and pant;
- A hammer and chisel;
- A pruning saw;
- Water spray(use where plant specimens are to be kept alive);
- Sturdy plant press, herbarium press and newspaper.

For entomologists:

- Sweep net;
- Pooter or aspirator;
- Lures or traps;
- Mounting boards and pins for insects;
- Cotton wool to place in tube with live insect (to prevent damage in field).

For plant pathologists:

- Spade and sieve for nematodes;
- Razor blades and scalpels (to section plant material);
- Culture plates;
- Specimen pots;
- Calcium chloride chips (to act as a desiccant);
- Ethanol;
- Ethanol flame lamp to sterilize scalpels, tweezers etc.

Checking for consistency in diagnostic skills of surveillance team members

To assess if members in the team observe and record pests in uniform manner, begin by selecting five or more infested plants and number them. Each team member then assesses all the plants, recording details per plant on their own. Compare the results between team members, both per plant and as an average over the five (or more) plants. If there are differences in the records, inspect the plants together to develop a consensus on the results. Repeat the process with new plants or sites until consistent results are obtained within the group. If there is debate concerning the diagnostic characteristics, seek further information about their appearance for the given conditions.

Obtain permits

Consider whether a permit is needed to visit villages, communities, or forests where it is intended to survey. It is essential to inform and involve people, as appropriate, particularly those in charge of the area; for example, CDO in the district. It is important to inform them on what dates the survey team would like to visit and give them a clear explanation of what the team will be doing and the expected results. The timing may coincide with cultural or ritual events and so access may be denied. It is wise to re-confirm permission before leaving. It may need to obtain release and transport permits from District Forest Office (DFO) and also quarantine permits for domestic and international transfer of specimens collected.

O. PERFORM THE SURVEY

The team should now be equipped with enough plans, information and tools to carry out the survey.

P. ANALYSE DATA

After survey, the surveyor should have a set of forms or data that are not processed or analyzed. The raw data can be used to:

- calculate basic statistics, such as the average and total numbers of pest;
- create a map of the pest distribution or spread;
- Create pest records and the pest list of the surveyed host in the surveyed area. The standard format that can be used for pest record is given in the annex 3.

Q. REPORT THE RESULTS

If the survey is designed for trade-related and quarantine purposes, NPPO seeks for soft and hard copy of the report before acceptance.

Summary report

It is useful to have a simple summary to be provided as follow-up information for the people who were involved in the survey; from the team members to the local property owners, rangers and community leaders. This acknowledges their assistance and show that their involvements are appreciated. Summary report should include the following sub-titles:

- the survey title and team members;
- the aim of the survey, including which pests, hosts and sites were targeted and why;
- What was found?
- What it means to the people who read the report?

Final report to NPPO

It may be obligatory to submit the information according to the format and content of pest reports covered in NSPM 13 and 17 particularly, for situations involving trading partners. During the assessment and acceptance of the survey plan, the NPPO provides the format (if needed additionally) according to the objective of the survey. But in general condition, the report should provide at least the following information:

- The survey title and team members;
- The reason for surveying;
- Background information on the pest, host and sites surveyed, including discussion of any earlier, related surveys;
- The survey design methods in detail, including site selection, timing of the survey, the type of data and specimens collected;
- Process of data analysis and interpretation;
- Conclusions of survey findings and relation of finding with the objective. The report should provide the filled up information in the annexes 1, 2 and 3.
- Literature citation, if applicable, must be provided in the body text and at the “references” heading at the end of the report.
- The report should also have a brief abstract at the beginning and should include a glossary of terms and acknowledgments (such as from whom permission and funding were received).

Special note on the difference between survey of cropping ecosystem and natural ecosystem

Natural ecosystem	Cropping ecosystem
Blitz survey preferred because of random distribution of host.	Randomization within the single unit preferred, because of the uniformity of the host distribution
Data transformation from the mean value to the designed confidence is not recommended, because of higher chance of sampling error in randomized survey.	Data transformation from the mean prevalence value to the designed confidence level is preferred.
Prior to conduct the pest survey in the natural ecosystem, it is always essential to get the report of host distribution survey.	Perceived Cropping pattern of an area will give the information on host distribution.
Difficulty of designing pest information system.	Possibility of designing farmer led pest information system is high.

Natural ecosystem	Cropping ecosystem
Early detection is very difficult, because of lower flow of the people in the natural ecosystem.	Early detection is easy, because of high flow of people in the agriculture ecosystem.
Incursion management of the invasive pests is easier because of limited chance of artificial spread	Incursion management is difficult because of high chance of artificial spread
Incursion is difficult because of the natural balance	Incursion is easier because of disturbed ecosystem.

SELECTED REFERENCES

- Binns, M.R., Nyrop, J.P. and van der Werf, W. 2000. *Sampling and monitoring in crop protection*. The theoretical basis for developing practical decision guides. CAB International, Oxon, UK and New York, USA.
- ISPM No. 5. 1997. *Glossary of phytosanitary terms*. FAO, Rome.
- ISPM 6. 1997. *Guidelines for surveillance*. IPPC, FAO, Rome.
- ISPM 8. *Determination of pest status in an area*. IPPC, FAO, Rome.
- NICRA team of rice pest surveillance. 2011. *Manual for Rice Pest Surveillance*. Jointly published by National Centre for Integrated Pest Management, New Delhi, Central Research Institute for Dryland Agriculture and Directorate of Rice Research, Hyderabad. 40 pp.
- Pokharel, S. 2009. *Training report on pest survey and surveillance & monitoring-I*, 16-23 March 2009, Plant Protection Directorate, MTF/NEP/060/STF (STDF-170)
- Report of the global symposium on plant pest surveillance, 29 October – 2 November, 2012, Anyang, Seoul, Republic of Korea
- Report of the regional Asia-pacific workshop for the global review of phytosanitary surveillance in the context of the IPPC standard (ISPM6), Identification of challenges and best practice, 31 January – 3 February 2012, Chiang Rai, Thailand
- ISPM 4. 1996. *Requirements for the establishment of pest free areas*. FAO, Rome.
- ISPM 10. *Requirements for the Establishment of Pest Free Places of Production and Pest Free Production Sites*.
- NPPO. 2014. *Technical guideline for detection survey of plant pests in Nepal* (Draft). Endorsed by NPPO of Nepal on 22 Baisakh, 2071, accepted by Plant Quarantine Committed on 16 Jesth, 2071.
- Teresa McMaugh, 2005. *Guidelines for surveillance for plant pests in Asia and the Pacific*, ICIAR, Govt. of Australia
- Various presentations of Dr. BN Mahato, and Dr. SL Joshi on pest survey training conducted at PPD and at NPQP in different fiscal years of the GON.
Guidelines for surveillance for plant pests in Asia and the Pacific, 2005
-will manage the reference and citation accordingly

अनुसूची-१

Pest Information Collection Format for National Pest Database of Nepal Generalized survey questionnaire

A. GENERAL INFORMATION

Title of the study/survey	
Name of field/site visited	
Date/time of field visit when the pest was intercepted	
Name and the address of local people involved	
Contact details of local people/s involved in the survey	Phone: Email:
GPS reference point	Latitude:
	Longitude:
	Altitude:
Locality	Village name: ward no.:
	Local level:
	District:
Climate data of locality	Average min. temp (in °C):
	Average max. temp (in °C):
	Rainfall (in mm)

B. DESCRIPTION OF HABITAT

Aspect (River bank/forest line/others),

Slope (north west/south west others),

Vegetation type phylogenetic options including keystone species (*Schima/Catonopsis*)

Soil type (refer 12 soil classification category triangle for example sand/silt/clay and their combination)

C. DESCRIPTION OF HOST

Scientific name of host:

Common name:

Family:

Variety:

Growth habit: (for example: herb/shrub/ tree/ vine/ succulent/ woody/ trailing etc.)

Habitat characters: wild/cultivated/protected/aquatic/arboreal/terrestrial/amphibian like ferns/public place/garden etc.....)

D. PHENOLOGICAL STAGE OF HOST SURVEYED

- Plant part/s affected
- First appearance of the pest (in which part/s of the plant):
Area affected by the pest (among total area of plants/crops):
Percent crop/plant affected quantified (10/20/30.....)
- Yield loss percent
- Major symptoms observed
- Distribution of symptoms within field
- Describe other problems observed (draught/nutritional/water logging/ biotic)
- Type of organism/s seen
- (Fungi, Bacteria, Insect/Mites, Nematode, Virus, Phytoplasma, Weed, Unknown)

E. DETAIL INFORMATION

- Life stage of pest observed/analyzed/diagnosed: [] Egg; [] Larvae (caterpillar/grub/maggot); [] Pupae; [] Nymph; [] Adult, anamorph, teleomorph, juvenile, dormant spore.....
- Life stage of pest affecting alternate/collateral host: (Farmers' view) [] Egg; [] Larvae (caterpillar/grub/maggot); [] Pupae; [] Nymph; [] Adult , anamorph, teleomorph (regular, irregular, rare, unknown), juvenile, dormant spore, etc (Name the host, if further survey in the field is required, make recommendation)
- Winter or summer diapause and relevant climatic cues; physiological adaptations for survival of low temperatures, desiccation etc. in or out of diapause (yes/no)
- Long term survival stage seen (eg. sclerotia/talia ...) in soil/ on cultivated hosts/ on wild hosts/ on obligate alternate hosts/on seeds/ on contaminated surfaces / machinery)
- Presence of other suitable plants that could be new hosts (yes/no), if further survey in the field is required, make recommendation
- Any alternate or collateral host observed around the field? (yes/no)
- Proximity of pest to other suitable hosts/habitats (indicate distance in meter-consider if the isolation distance is managed in case of seed production fields around)
- Abundance of main hosts and alternate hosts and how they are distributed (dense/sparse)-consider the isolation distance for seed production to judge, if further survey in the field is required, make recommendation
- Any vector seen? (Name the vector, if further survey in the field is required, make

recommendation)

- If transmitted by vectors, if the suitable vectors are available in the survey area (yes/no), if further survey in the field is required, make recommendation
- Duration of the damaging life stage as observed by the farmers (in days/weeks)
- Other seasonal timing of pest appearance as observed by the farmers (Spring/Summer/Rainy/Autumn/Winter)
- Dispersal mechanism of pest as observed by the farmers (air/water/soil/plant/animal/insect)
- Destination of the produce (domestic / international) (Name the place/s in the separate comment box)
- Use of by-products and waste (feed/fuel/manure/fodder/litter/other)
- Soil factor affecting pest survival (acidic/alkaline/upland/lowland/fertile/marginal.....)
- Irrigation practice affecting pest survival (furrow/flood/check basin/drip/sprinkler.....)
- Planting method affecting pest survival (continuous row sowing/broadcasted/transplanted.....)
- Crop geometry affecting pest survival (recommended/non-recommended.....)
- Natural enemies observed (yes/no)
- If natural enemies observed name those (if further survey in the field is required, make recommendation in the separate box)
- Abundance of natural enemies (dense/sparse), if further survey in the field is required, make recommendation in the separate box)
- Natural barriers around affecting spread of pest (river/lake/forest/mountain/city) if further survey in the field is required, make recommendation in the separate box)
- Any qualitative effect significantly observed due to particular pest (Effect in community structure/Physical modifications of habitats/Changes in nutrient cycling and availability/Changes in trophic and mutualistic interactions/Unemployment/Health effects on human and animals/Recreation/tourism/education/spiritual impacts/Impact on aesthetics (if further survey in the field is required, make recommendation)
- Number of generations per year: univoltine/multivoltine (1/2/3/4/5/6/)
- parthenogenetic multiplication (yes/no)
- possible number of infection cycles per growing season: polyictic/multiectic (1/2/3/4/5/6)
- Pest management practice of the local people
- Pest management recommendations from the surveyor

F. DIAGNOSTIC INFORMATION

Scientific name of pest:

Common name:

Growth habit:

Habitat characters:

Species name:

Family:

Order:

Subspecies, pathotypes, *formaespeciales*, overlapping species, synonymy

- Method of identification: (morphological/taxonomic/molecular)
- Name of the pest collector:
- Name and address of the diagnostic laboratory
- Name and organization affiliation of the identifier:
- Contact details of identifier: Phone: _____ Email: _____
- Date of verification:
- Method of verification: (morphological/taxonomic/molecular...)
- Name and organization affiliation of the verifier:
- Name and address of verification laboratory:
- Method of pest preservation (herbarium/culture/others)
- Habit character of pest
- Preserved item suitable for: Visual/Taxonomic/molecular analysis
- Name and address of pest library (Laboratory) preserving the pest
- Life stage of pest on the preserved sample [] Egg; [] Larvae (caterpillar/grub/maggot); [] Pupae; [] Nymph; [] Adult, anamorph/ teleomorph/ juvenile/dormant spore/sclerotia/refined genetic material)
- Is this a quarantine pest? (yes/no)
- Phylogenetic relationship with any known quarantine species? (yes/no)
- Further study recommended?
- Species accumulation curve: satisfied consecutive sites /not-satisfied
- Further study recommended?
- Recommended for RNQP (yes/no) or further survey required? (yes/no)
- Further study recommended?
- Any additional information (including collection of specimens for investigation):

Name, signature/date of surveyor/s:

अनुसूची-२

FORMAT FOR SPECIMEN FORWARDAL

Specimen forwarded for identification by Diagnostic/Referral Laboratory	
1. Collection number:	2. Date of Collection:
3. Submitting organization:	
4. Name/Address of the sender:	
5. Locality of collection (District/local level/village):	
6. Reasons for identification:	
7. Name of the host plant species (Common/Scientific)/variety and or/ commodity:	
8. Origin of host/commodity (where applicable):	
9. Plant parts affected: <i>*tick out in appropriate box</i>	<input type="checkbox"/> roots; <input type="checkbox"/> stems; <input type="checkbox"/> leaves; <input type="checkbox"/> inflorescence; <input type="checkbox"/> fruits; <input type="checkbox"/> seeds/nuts <input type="checkbox"/> others (_____)
10. Category of pest specimen/organism submitted: <i>*tick out in appropriate box</i>	<input type="checkbox"/> insects; <input type="checkbox"/> mites; <input type="checkbox"/> nematodes; <input type="checkbox"/> fungi; <input type="checkbox"/> bacteria; <input type="checkbox"/> virus; <input type="checkbox"/> others (_____)
11. Life stage of the pest (Applicable to insects): <i>*tick out in appropriate box</i>	<input type="checkbox"/> egg; <input type="checkbox"/> larvae; <input type="checkbox"/> pupae; <input type="checkbox"/> adult; <input type="checkbox"/> nymphs; <input type="checkbox"/> juveniles; <input type="checkbox"/> anamorphic <input type="checkbox"/> ; cysts; <input type="checkbox"/> others (_____)
12. Type of pest specimen/organism submitted: <i>*tick out in appropriate box</i>	<input type="checkbox"/> preserved specimen; <input type="checkbox"/> pinned/card board mounted specimen; <input type="checkbox"/> dry specimen with host; <input type="checkbox"/> culture; <input type="checkbox"/> disease specimen (fresh); <input type="checkbox"/> disease specimen (partially dry); <input type="checkbox"/> slide mount; <input type="checkbox"/> others (_____)
14. Number of specimens submitted per each collection:	

15. Signature/stamp/office seal of the sender with date: _____	
For identifier use	
16. Name & address of Diagnostic/Referral Laboratory:	
17. Remarks of identifier (condition of receipt of specimens):	
18. Pest identification (Common/Scientific name/Taxon):	
19. Description notes, if any:	
Place: _____	
Date: _____	

(Signature/Name/Designation of Identifier)	
Note: This form should be prepared in duplicate by the sender and forwarded to the identifier/referral laboratory along with each collection of specimen. The identifier should return the original copy after entering the particulars of the pest identified along with description notes and remarks if the identifier will retain any to the sender of the specimen and duplicate copy.	

**PEST INFORMATION REPORTING FORMAT FOR NATIONAL PEST
DATABASE OF NEPAL**

Abstract

A. Background information

1. Introduction

- The survey title and
- Survey team member's/specimen collector's name:
- Background information on the pest, host and sites surveyed, including discussions of any earlier, related surveys:
- The survey design/methods/protocols, including site selection and sampling, timing of the survey, the type of the data and specimen collected:
- General findings
- Process of data analysis and interpretation:
- Survey protocol applied (Methodology):
- Define the scope of survey
 - Reason for Survey
 - Status of previous survey for this particular pest
 - Possible cause of absence or presence of pest in the survey area
- Individual Pest Information

1. Pest Identity:

- Provide details on taxonomy and nomenclature
- Pest diagnostic protocol, diagnostic/verifying laboratories
- Pest diagnostician

2. Regulatory status of the pest

- Define if the pest is already regulated or has been previously assessed for the country/area at risk:
- Quarantine pest absent from the country
- QP present in the country, but not widely distributed and under official control
- RNQP but whose presence in the plants for planting affects its intended use intensely
- Pest requiring phytosanitary measure (not officially listed)

3. Regulatory status of the pest elsewhere

- Provide details of the pest's regulatory status in other areas/countries

4. **Distribution summary**
 - Provide a summary of the global distribution of the pest, the pest's presence in neighboring countries, areas with similar climates and countries with strong trade and transport links to the survey area
5. **Association with host plants or habitats**
 - Describe host plants, or suitable habitats, that occur in the survey area (outdoors, in protected cultivation or both)
6. **Potential conditions for pest in the survey area**
 - Describe eco-climatic conditions area that is supporting the establishment and spread of the pest (including presence of alternate hosts and vectors where relevant; consider protected conditions as well)
7. **Economic, social and environmental impact in the survey area**
 - Provide evidence that the pest has potential to have unacceptable economic, environmental and social impact in the survey area
8. **The detected pathways of pest association in specific commodity**
 - Plants for planting
 - Budwood/graftwood
 - Bulb tuber /corm/rhizome for planting
 - Seeds for planting
 - Tissue culture
 - Plants for planting
 - Not for planting
 - Bark
 - Cut flower/branches
 - Dried or stored plant products
 - Fresh fruit or vegetable
 - Grain/seed
 - Non squared wood
 - Pollen/honey
 - Squared wood
 - Transport stowaway
 - Aircraft
 - Container or bulk
 - Containers and packaging - non wood
 - Containers and packaging - wood
 - Land vehicle
 - Machinery/equipment
 - Manufactured goods
 - People and their luggage/equipment
 - Ship/boat

- Sand soil gravel compost
- Unaided
 - Water
 - Wind
 - Other ways

B. Specific description (Please provide quantitative data where relevant)

- Occurrence of life stage of pest associated with pathway/commodity
- Seasonal timing of occurrence of pest in the survey area
- Pest management procedures applied by the local people and its effect in product output
- Duration and vulnerability of life cycle of pest
- Previous occurrence of the pest
- Local procedures for pest freedom in the final product
- Field inspection methods and quality control of the product
- Certification schemes applied by the farm (if any)
- Chemical treatment options available in the field
- Dispersal mechanisms of host and pest, including vectors
- Number and name of destinations of the final product
- Proximity of pest to other suitable hosts/habitats
- Intended use of the commodity including by-products and waste
- Abundance of main hosts and alternate hosts and how they are distributed
- Presence of other suitable plants that could be new hosts
- If transmitted by vectors, explain if the suitable vectors are available in the survey area
- Explain the observed soil factors for soil borne pests
- Cultural practices e.g. irrigation, planting, harvesting methods that affects pest survival and spread etc.
- If observed any existing natural enemies, explain those.
- Reproductive and survival strategies of pest (Literature review)
- Genetic adaptability (Literature review)
- Climate change factors that affect the pest's reproduction and survival
- Climate change factors that affect the pest's hosts, vectors or habitats
- Natural barriers for the pest spread in the survey area

- Number of generations of pest and host per year

C. **Consequences observed:**

Quantitative data

1. Reduction in crop yield or quality
2. Reduction in prices or demand, including export markets
3. Increase in production costs (including costs of control)
4. Threat to other local species, with special focus on threatened and keystone species
5. Hybridization with native species (Literature review on variability)

Qualitative data

1. Effect in community structure
2. Physical modifications of habitats
3. Changes in nutrient cycling and availability
4. Changes in trophic and mutualistic interactions
5. Unemployment
6. Health effects on human and animals
7. Recreation, tourism, education or spiritual impacts
8. Impacts on aesthetics

D. **Conclusions and Recommendations**

Note: Literature citation, if applicable, must be provided in the body text and at the references heading at the end of the report (If the data is extracted from plant clinics or from the IPM field school please provide the photocopy of record sheet). The report should also have a brief abstract at the beginning and should include high resolution photographs, address or voucher specimen library, a glossary of terms and acknowledgements (such as from whom permission and findings were received).

PART -2

WORK FLOW FORMATS FOR PRA

फाराम नं.-१

INFORMATION REQUIRED FOR PEST RISK ANALYSIS FOR IMPORT OF PLANT AND PLANT MATERIALS INTO NEPAL

(To be provided by the prospective EP applicant while requesting or recommending for the EP, Pursuant to Plant Quarantine and Protection Rule 2066 (5))

1. Details of Applicant

1.1 Name/ Organisation.....

1.2 Address.....

1.3 Contact details..... E-mail.....

2. PRA General Parameters

2.1 Scientific & Common name of the product.....

2.2 Country of origin.....

2.3 Quantity/ Volume to be imported

3. Product Type (please mark one or more)

3.1 Processed/ Non-processed

3.2 Living/ dead

3.3 Plant/ Animal (including beneficial insects)

3.4 Genetically modified/ non-genetically modified

3.5 Seed/ sapling (if sapling, with soil or soilless)

3.6 Culture / non-culture

3.7 other (please specify).....

4 Product Processing (if applicable)

4.1 If seed: ground/ kibbled/ whole/ preserved

4.2 If plant: fresh/ dried/ freeze dried/ preserved

4.3 Processing refinement: cooked/ frozen/ pulped/ steamed

4.4 Specify treatment details

5. Product Origins (please state if question not relevant)

5.1 Source location (by country, origin & locality)

5.2 Production method, Certification scheme and / or accreditation type?

.....

6 End Use (circle one or more)

6.1 Human consumption / Processing/ Stock feed/ Pet food/ Fish food/ Seeds for sowing/
Nursery stock/ Multiplication/ pre-entry Quarantine studies/ Therapeutic/ Fertilizers/ In-vivo /
In-vitro

6.2 Other

7 End Destination (circle &/or specify)

7.1 Rural/ urban

7.2 Multiple locations/ single

7.3 Specify Country, State & / or region (PRA defined area)

8. Entry mode (circle one or more): Ship/ Air/ Ground transport /rail/other.....

9. General Comments (any further general comment or notes that need to be made, please
make here).....

INFORMATION NEEDED TO UNDERTAKE PRA IN NEPAL

(To be provided by the exporting country)

1. Plant and Plant Product Intended to Export in Nepal

- 1.1 Common name
- 1.2 Scientific (genus & species/strain/variety/cultivar) name
- 1.3 Resistant or non-resistant varieties (Provide the name of the pests and the corresponding resistant/non-resistant varieties)
- 1.4 Other countries that have already imported the proposed product
- 1.5 Plant part to be imported by Nepal (whole plant/seed/cutting/sapling/ budwood/ bulb/fruit/grafted plants)

2.2 Production and Export (tons/year);

3. Cultivation practices

4. Packaging

- 4.1 Method of packaging
- 4.2 Inspection procedure
- 4.3 Post harvest treatment
- 4.4 Conditions and security of storage place

5. Export program (policy/activity)

- 5.1 other trading partners
- 5.2 Existing procedure for issuing phytosanitary certificates (including additional declaration).

6. Pest List (separately for all the pests)

6.1. Pest Identity:

- a. Provide details on taxonomy and nomenclature
- b. Generally, the taxonomic unit of the pest is the species; the use of higher or lower taxonomic units should be supported by a scientifically sound rationale

- 1.2. Survey protocol applied
- 1.3. Regulatory status of the pest
- 1.4. Provide a summary of the distribution of the pest.

Describe eco-climatic conditions of the area that is supporting the establishment and

spread of the pest (including presence of alternate hosts and vectors where relevant; consider protected conditions as well)

1.5. The detected pathways of pest association in specific commodity

1.6. Specific description

- Occurrence of life stage of pest associated with pathway/commodity
- Seasonal timing of occurrence of pest in the production area
- Duration and vulnerability of different stages of life cycle of pest
- Previous occurrence of the pest in the production area
- Local procedures for pest freedom in the final product
- Field inspection methods and quality control system of the product
- Certification schemes applied by the farm
- Chemical treatment options applied in the field
- Dispersal mechanisms of the pest, including vectors
- Number and name of destinations of the final product
- Intended use of the commodity including by-products and waste
- Collateral and alternate hosts available in the production area
- If transmitted by vectors, explain if the suitable vectors are available in the production area
- Cultural practices e.g. irrigation, planting, harvesting methods their relationship with the pest survival
- If observed any existing natural enemies, explain those.
- Number of generations of pest and host per year
- Reduction records in crop yield or quality

7. Relevant Documents References including URL access link:

FORMAT FOR PEST RISK ANALYSIS
RISK OF INTRODUCING PESTS VIA (Commodity type) FROM (Country) INTO
NEPAL

Crop(s): Scientific Name (Common name)

Commodity types(s):

PRA NO.:

Month and Year of PRA:

Expert's Reference Number:

GENERAL INFORMATION:

Initiation and scope of the PRA

PRA area

Do previous PRAs exist for this or a similar pathway?

Weed potential of the commodity

Description of the pathway

Why the import is necessary?

Risk Category:

Processing Step After Entry and Processing Destination

National Annual Production Status

Existing Marketing Problem of Local Product

National Annual Consumption Pattern

Volume/quantity of commodity required to be imported?

RAPID PEST RISK ASSESSMENT

APPENDIX 1. Full pest list for the comparison between (china, for example) and Nepal against (cowpea seed, for example) import.

Type	Species	On crop	On commodity type Stage-1 (S1)	On commodity type Stage-3 (S3)	Nature of association on commodity type	Country/area of origin	Importing country/area Nepal	Included in the assessment?	Source of pest record

APPENDIX 2. Pests potentially requiring phytosanitary measures

Type	Species	RS, Stage-1 (S1)	RS, S2	RS, S3	RS, S4	RQ1	RQ2	RQ3	RQ4	RQ5	References

Regulatory status (RS)

- a. No regulation required/ Unknown (Appendix 4)
- b. Quarantine pest, absent from the country/area (Detail in Appendix 3 C)
- c. Quarantine pest, present in the country/area but not widely distributed and under official control (Appendix 3C)
- d. Regulated non-quarantine pest, present in the country/area but whose presence in plants for planting affects its intended use (Appendix 3 D)
- e. Pest requiring phytosanitary measures (not yet officially listed) (Appendix 3D)
- f. Pests for deregulation (Appendix 4 A/RS5)
- g. Predators/parasitoids/antagonists/bio-control agents (Appendix 5/RS6, if any)
- h. Vectors of RS1/RS4 pests (Appendix 6, RS7)
- i. National pest of Crop (Appendix 3A)
- j. National pest of Commodity Type (Appendix 3B)

RAPID ANSWERS

Pests	RS	RQ1: Does this pest have potential for establishment and spread in the PRA area?	RQ2: Does this pest have potential for economic, environmental and social consequences in the PRA area?	RQ3: Does this pest requires PEQ surveillance? What is the duration of PEQ tracking?	RQ 4: Vector of QP/PQP?	RQ5: Potentially hyper parasite?	Proposed RS

APPENDIX 3. Pest categorization

A. National pest list of (black pepper, for example) that may not appear on commodity types of trade

Type	Species	On crop	On commodity type (Stage 1)	On Commodity type (Stage 3 from Appendix 1)	Source of pest record	In country surveillance and survey need

B. National pest list of (black pepper, for example) that may appear in the commodity type of trade while exporting phytosanitary certificate from Nepal may/could/should declare the freedom of these pests (according to the requirement of importing country)

Type	Species (Including Added ones from Stage 3 Appendix 1)	On crop	On commodity type	Source of pest record	In country surveillance and survey need

C. Quarantine pests

Type	Species	Nature of association with commodity type	RS	Source of pest record	Risk Assessment/ Management	Pest requires phytosanitary measures?

D. Potential quarantine pests

Type	Species	Nature of association with commodity type	RS	Proposed RS	Source of pest record	Risk Assessment for surveillance and management instruction	Pest requires phytosanitary measures?

APPENDIX 4: Pests for deregulation (RS5)

Type	Species	Nature of association with commodity type	Current RS	Proposed RS	Source of pest record	Risk Assessment for in country surveillance and management instruction	Pest requires phytosanitary measures?

APPENDIX 5: Predators/Parasitoids/Antagonists/Bio-control agents/ beneficial insects (/RS6, IF ANY)

Type	Species	Nature of association with commodity type	RS	Source of record	Assessment for procurement and use	Requires phytosanitary measures?

Appendix 6: Vectors of RS1/RS4 pests (RS7)

Type	Species	Nature of association with commodity type	RS	Source of record	Assessment of risk	Requires phytosanitary measures?

- Recommendation of rapid PRA
- List of pests recommended for full PRA (fPRA)

FULL PEST RISK ANALYSIS (FOR EACH PEST OF APPENDIX 3D)

Name and Contact Details of the Risk Analysts

Pest name:

Pest Identity:

Provide details on taxonomy and nomenclature

Generally, the taxonomic unit of the pest is the species; the use of higher or lower taxonomic units should be supported by a scientifically sound rationale

.....

1. Risk assessment

1.1. Probability of entry

1.1.1. How might the pest enter the PRA area? Recognize relevant pathways for introduction. Pathways can represent any means that allow the entry or spread of a pest. Selected pathways will be assessed individually for probability of entry and risk management.

1.1.2. What is the probability of the pest being associated with the pathway at origin?

Factors to consider

- Prevalence of pest in the source area
- Occurrence of life stage able to associate with pathway
- Seasonal timing
- Pest management procedures applied at place of origin

Rating	Judgment	Confidence Level	Judgment
Non Assessed		Non Assessed	
Negligible		Negligible	
Low		Low	
Medium		Medium	
High		High	

1.1.3. What is the probability that the expected volume and frequency of import of the commodity will support entry?

Rating: Low

Confidence: High

1.1.4. What is the probability of the pest surviving during transport?

Factors to consider

- Speed and conditions of transport
- Duration and vulnerability of life cycle
- Previous interceptions of the pest
- Commercial procedures during transport (e.g. refrigeration)

Rating	Judgment	Confidence Level	Judgment
Non Assessed		Non Assessed	
Negligible		Negligible	
Low		Low	
Medium		Medium	
High		High	

1.1.5. What is the probability of the pest surviving or evading existing pest management procedures?

Factors to consider

- Inspection methods and quality control
- Certification schemes
- Chemical treatment

Rating	Judgment	Confidence Level	Judgment
Non Assessed		Non Assessed	
Negligible		Negligible	
Low		Low	
Medium		Medium	
High		High	

1.1.6. What is the probability of transfer to a suitable host or, in the case of potential weeds, habitat?

Factors to consider

- Dispersal mechanisms, including vectors
- Dumber of destinations
- Proximity to suitable hosts/habitats
- Seasonality
- Intended use of the commodity, in the case of a commodity pathway (e.g. for planting, processing, consumption)
- Risks from by-products and waste

Rating	Judgment	Confidence Level	Judgment
Non Assessed		Non Assessed	
Negligible		Negligible	
Low		Low	
Medium		Medium	
High		High	

1.1.7. Summary of rating and summary of confidence level of above 5 questions

- Do you consider this pathway a major or a minor pathway...
.....
.....
- Does this pathway require management measures... ..
.....

Summary on probability of entry

Pathway	Summary rating	Summary confidence level	Major/ minor pathway?	Pathway requires management measures?

1.1.8. References:

1.2. Probability of establishment

1.2.1. What is the probability that suitable hosts or, in the case of potential weeds, habitats are available in the PRA area?

Factors to consider

- Abundance of main hosts and alternate hosts and how they are distributed
- Geographic proximity of hosts/habitats to pathway destinations
- Presence of other suitable plants that could be new hosts

Rating	Judgement	Confidence Level	Judgement
Non Assessed		Non Assessed	
Negligible		Negligible	
Low		Low	
Medium		Medium	
High		High	

1.2.2. If transmitted by vectors, what is the probability that suitable vectors are available in the PRA area?

Rating	Judgement	Confidence Level	Judgement
Non Assessed		Non Assessed	
Negligible		Negligible	
Low		Low	
Medium		Medium	
High		High	

1.2.3. What is the probability that climatic conditions and other abiotic factors will allow the pest to establish in the PRA area?**Factors to consider**

- Compare the known distribution of the pest with eco-climatic zones in the PRA area
- Whether hosts are grown in protected cultivation
- Soil factors for soil-borne pests

Rating	Judgement	Confidence Level	Judgement
Non Assessed		Non Assessed	
Negligible		Negligible	
Low		Low	
Medium		Medium	
High		High	

1.2.4. What is the probability that existing control measures for other pests in the PRA area are unable to prevent establishment?

Factors to consider

- Cultural practices e.g. irrigation, planting, harvesting methods etc.
- Pest control programs

Rating	Judgement	Confidence Level	Judgement
Non Assessed		Non Assessed	
Negligible		Negligible	
Low		Low	
Medium		Medium	
High		High	

1.2.5. What is the probability that existing natural enemies in the PRA area are unable to prevent establishment?

Rating	Judgement	Confidence Level	Judgement
Non Assessed		Non Assessed	
Negligible		Negligible	
Low		Low	
Medium		Medium	
High		High	

1.2.6. What is the probability that other biological characteristics of the pest will enable establishment?

Factors to consider

- Reproductive and survival strategies
- Genetic adaptability
- Minimum population needed for establishment

Rating	Judgement	Confidence Level	Judgement
Non Assessed		Non Assessed	
Negligible		Negligible	
Low		Low	
Medium		Medium	
High		High	

1.2.7. What is the probability of establishment under foreseeable climate change conditions?

Factors to consider

- Climate change projection
- Climate change factors that affect the pest’s reproduction and survival
- Climate change factors that affect the pest’s hosts, vectors or habitats

Rating	Judgement	Confidence Level	Judgement
Non Assessed		Non Assessed	
Negligible		Negligible	
Low		Low	
Medium		Medium	
High		High	

1.2.8. Summary of probability of establishment

- Summary of rating and confidence level
.....
- Summary note on establishment.....
.....

1.2.9. References

1.3 Probability of spread

1.3.1 What is the expected rate of natural spread in the PRA area?

Factors to consider

- Rate and distance of spread elsewhere
- Natural barriers in PRA area

Rating	Judgement	Confidence Level	Judgement
Non Assessed		Non Assessed	
Negligible		Negligible	
Low		Low	
Medium		Medium	
High		High	

1.3.2. If transmitted by vectors, what is the expected rate of spread by vectors in the PRA area?

Factors to consider

- Rate and distance of spread elsewhere

Rating	Judgement	Confidence Level	Judgement
Non Assessed		Non Assessed	
Negligible		Negligible	
Low		Low	
Medium		Medium	
High		High	

1.3.3. What is the expected rate of spread with commodities or conveyances in the PRA area?

Rating	Judgement	Confidence Level	Judgement
Non Assessed		Non Assessed	
Negligible		Negligible	
Low		Low	
Medium		Medium	
High		High	

1.3.4. What is the probability of the pest spreading to an area of higher economic importance than the area of introduction?

Rating	Judgement	Confidence Level	Judgement
Non Assessed		Non Assessed	
Negligible		Negligible	
Low		Low	
Medium		Medium	
High		High	

1.3.5. If a commodity pathway has been identified as one of the pathways of entry, what is the probability that the intended use of the commodity increases the rate of spread?

Factors to consider

- Whether intended for planting, processing or consumption
- Disposal of waste, by-products
- Number and location of expected destinations

Rating	Judgement	Confidence Level	Judgement
Non Assessed		Non Assessed	
Negligible		Negligible	
Low		Low	
Medium		Medium	
High		High	

1.3.6. What is the potential rate of spread under foreseeable climate change conditions?

Factors to consider

- Climate change projection
- Climate change factors that affect the dispersal of the pest

Rating	Judgement	Confidence Level	Judgement
Non Assessed		Non Assessed	
Negligible		Negligible	
Low		Low	
Medium		Medium	
High		High	

1.3.7. Summary of probability of spread

- Summary of rate and Confidence level.....
- Summary note on spread.....

1.3.8. References

1.4 Potential consequences

1.4.1. What is the level of economic loss to agriculture, forestry or horticulture associated with this pest in its existing geographic range?

Factors to consider

- Reduction in crop yield or quality
- Reduction in prices or demand, including export markets
- Increase in production costs (including costs of control)
- Vectoring of other pests of economic importance

Rating	Judgement	Confidence Level	Judgement
Non Assessed		Non Assessed	
Negligible		Negligible	
Low		Low	
Medium		Medium	
High		High	

1.4.2. What is the level of potential economic loss to agriculture, forestry or horticulture in the PRA area?

Factors to consider

- Reduction in crop yield or quality
- Reduction in prices or demand, including export markets
- Increase in production costs (including costs of control)
- Vectoring of other pests of economic importance

Rating	Judgement	Confidence Level	Judgement
Non Assessed		Non Assessed	
Negligible		Negligible	
Low		Low	
Medium		Medium	
High		High	

1.4.3. What is the level of negative impact on native biodiversity associated with this pest in its existing geographic range?

Factors to consider

- Threat to native species, with special focus on threatened and keystone species
- Changed community structure
- Hybridization with native species

Rating	Judgement	Confidence Level	Judgement
Non Assessed		Non Assessed	
Negligible		Negligible	
Low		Low	
Medium		Medium	
High		High	

1.4.4. What is the level of potential negative impact on native biodiversity in the PRA area?

Factors to consider

- Threat to native species, with special focus on threatened and keystone species
- Changed community structure
- Hybridization with native species

Rating	Judgement	Confidence Level	Judgement
Non Assessed		Non Assessed	
Negligible		Negligible	
Low		Low	
Medium		Medium	
High		High	

1.4.5. What is the level of negative impact on ecosystem patterns and processes associated with this pest in its existing geographic range?

Factors to consider

- Physical modifications of habitats
- Changes in nutrient cycling and availability
- Modifications of natural successions
- Changes in trophic and mutualistic interactions

Rating	Judgement	Confidence Level	Judgement
Non Assessed		Non Assessed	
Negligible		Negligible	
Low		Low	
Medium		Medium	
High		High	

1.4.6. What is the level of potential negative impact on ecosystem patterns and processes in the PRA area?

Factors to consider

- Physical modifications of habitats
- Changes in nutrient cycling and availability
- Modifications of natural successions
- Changes in trophic and mutualistic interactions

Rating	Judgement	Confidence Level	Judgement
Non Assessed		Non Assessed	
Negligible		Negligible	
Low		Low	
Medium		Medium	
High		High	

1.4.7. What is the level of negative social impact associated with this pest in its existing geographic range?

Factors to consider

- Unemployment
- Health effects
- Recreation, tourism, education or spiritual impacts
- Aesthetics

Rating	Judgement	Confidence Level	Judgement
Non Assessed		Non Assessed	
Negligible		Negligible	
Low		Low	
Medium		Medium	
High		High	

1.4.8. What is the level of potential negative social impact in the PRA area?

Factors to consider

- Unemployment
- Health effects
- Recreation, tourism, education or spiritual impacts
- Aesthetics

Rating	Judgement	Confidence Level	Judgement
Non Assessed		Non Assessed	
Negligible		Negligible	
Low		Low	
Medium		Medium	
High		High	

1.4.9. What is the level of potential negative impact in the PRA area (for all sectors) under foreseeable climate change conditions?

Factors to consider

- Climate change projection
- Climate change factors that affect the pest’s reproduction and feeding habits
- Number of generations per year

Rating	Judgement	Confidence Level	Judgement
Non Assessed		Non Assessed	
Negligible		Negligible	
Low		Low	
Medium		Medium	
High		High	

1.4.10. Summary of potential consequence

	Summary Rating	Summary Confidence Level	Major/Minor Pathway?	Pathway Requires Management Measures?
Probability of entry (Pathway/s)				
Probability of Establishment				
Probability of Spread				
Potential Economic, Environmental and Social Consequences				

References

2. Risk Assessment Summary

- a. Does the pest require Phytosanitary Measure...?

3. Risk management

A. At the place of production

Management option	
Inspection or testing	
Post-harvest treatment	
Treatment of crop, field or place of production	
Specially protected growing conditions	
Maintenance of pest free area, place or site of production	
Resistant cultivars	
Specified harvest time to reduce crop infestation	
Certification scheme	
Other	

B. After harvest and during transport

Management option	Recommendations
Inspection or testing	
Post-harvest treatment	
Removal of specified parts of the plant	
Conditions for preparing and packing and storage	
In-transit conditions and treatment	
Restriction on period of entry (to reduce availability of susceptible hosts)	
Other	

C. At point of entry

Management option	Recommendations
Inspection or testing at point of entry	
Treatment at point of entry	

D. After entry

Management option	Recommendations
Inspection or testing in post-entry quarantine	
Surveillance, containment and eradication	
Restriction on end use or distribution	

E. Other

Management option	Recommendations
Prohibition of commodity	

Phytosanitary certification and additional declarations	
Risk management for this species is covered under other pest(s)	
No risk management option identified for this pest	

A. Instructions to quarantine inspectors at entry point

1. Pest name:

Symptoms (detection criteria):

Inspection Guide:

Action:

B. Instructions to the quarantine pest diagnostic laboratories at border offices

1. Lab Testing Methods for Quarantine Pests

A. Insect Pests

Pest	Testing Method	Detection Criteria	Action

B. Fungal Pathogens

Pathogen	Testing Method	Detection Criteria	Action

C. Bacterial Pathogens

Bacterium	Testing Method	Detection Criteria	Action

D. Phytoplasma Pathogen

Bacterium	Testing Method	Detection Criteria	Action

E. Nematodes

Nematodes	Testing Method	Detection Criteria	Action

C. REPORTING & COMPLIANCE

- **Positive Test:** If a quarantine pest is detected, the consignment must be **quarantined, re-exported, or destroyed.**
- **Negative Test:** If no pests are found, the consignment is cleared for **customs clearance.**
- **Documentation:** Lab results must be submitted to the **Nepal Plant Quarantine Authority** for final approval.

These **testing protocols** ensure imports comply with Nepal's **phytosanitary standards**, preventing the introduction of quarantine pests.

D. INSTRUCTIONS TO THE CENTRAL REFERAL SPS LABORATORY FOR DIAGNOSTICIANS

Molecular Primers for PCR Testing

1. Fungal Pathogens

Pathogen	Target Gene	Forward Primer (5' → 3')	Reverse Primer (5' → 3')	Amplicon Size (bp)	PCR Type

2. Bacterial Pathogens

Bacterium	Target Gene	Forward Primer (5' → 3')	Reverse Primer (5' → 3')	Amplicon Size (bp)	PCR Type

3. Phytoplasma

Phytoplasma	Target Gene	Forward Primer (5' → 3')	Reverse Primer (5' → 3')	Amplicon Size (bp)	PCR Type

4. Nematode

Nematodes	Target Gene	Forward Primer (5' → 3')	Reverse Primer (5' → 3')	Amplicon Size (bp)	PCR Type

These **molecular primer sets** allow precise **quarantine pest detection** for imports, ensuring compliance with Nepal's **phytosanitary regulation.**

E. INSTRUCTIONS TO THE NEPALESE IMPORTERS

- 1. Obtain Entry Permit from NPPO-Nepal before Everything:** Before entering into any trade agreement, finalizing payment terms (e.g., LC, TT, DAP, DAC, DAA, etc.), or engaging in any customs-related activities, it is mandatory to obtain an Entry Permit from NPPO-Nepal. This is a non-negotiable and critical requirement that must be fulfilled prior to initiating any import procedures for plants and plant-related products including walnut fruit requiring a Phytosanitary Certificate.
- 2. Phytosanitary Certificate Requirements:** The Phytosanitary Certificate issued by the NPPO-(Exporting country) must strictly adhere to Nepal's regulations and be issued only upon receipt of the Entry Permit from NPPO-Nepal. The Quarantine Authority in Nepal **shall not** put any requests for the release procedure of consignments that arrive at the entry point without meeting the required compliance. Traders are strongly urged to adhere to these regulations to avoid disruptions, penalties, and delays. The original Phytosanitary Certificate issued by NPPO-China must reference the **Approval Reference Number** from the **Entry Permit (EP)** issued by NPPO-Nepal.

Additional Declaration: The Phytosanitary Certificate (PC) must include an additional declaration confirming freedom from following pests, in strict adherence to the Entry Permit (EP) conditions issued by the National Plant Protection Organization of Nepal (NPPO-Nepal).

- 3. Pre-Shipment Fumigation:**

The consignment must undergo mandatory fumigation treatment before shipment to Nepal. The fumigation process must comply with international quarantine standards and must be certified by the NPPO of the exporting country. The fumigation details, including the treatment date, chemicals used, dosage, and exposure period, must be explicitly stated in the Phytosanitary Certificate.
- 4. Inspection Upon Arrival:**

Upon arrival at the designated port of entry in Nepal, all consignments will be subject to a mandatory phytosanitary inspection by the Quarantine Authority of NPPO-Nepal. The inspection will verify compliance with the Entry Permit conditions, Phytosanitary Certificate declarations, and freedom from the listed quarantine pests. Consignments failing to meet these requirements may be detained, treated (if feasible), returned to the country of origin, or destroyed at the importer's expense, as determined by NPPO-Nepal.

5. Packaging & Contamination-Free Requirement:

The garlic must be packed in clean, pest-free, and tamper-proof packaging to prevent any risk of contamination during transit. Packaging materials should comply with international phytosanitary standards and must not contain soil, plant debris, or any unauthorized organic matter. The consignment must be clearly labeled with product details, country of origin, and quarantine treatment information.

6. Approved Ports of Entry:

7. Importer Responsibilities:

- Hold importers accountable for ensuring that their consignments comply with Nepal's phytosanitary regulations, including bearing the costs of inspection, post entry confinement in the warehouse or the advanced diagnostic service. Implementing these measures would help Nepal safeguard its agricultural sector from potential threats posed by quarantine pests associated with imports.

फाराम नं. ४

ALOP FORM

नेपाल सरकार

कृषि तथा पशुपन्छी विकास मन्त्रालय

प्लान्ट क्वारेन्टाइन तथा बिषादी व्यवस्थापन केन्द्र

हरिहरभवन, ललितपुर

Appropriate level of protection against introducing (commodity type) from (country) into Nepal (for example)

(To be agreed and uploaded by the importer while applying for Entry Permit)

Entry Permit OF COMMODITY TYPE from NPPO-Nepal kindly invites the additional declaration of the freedom of following pest via the original PC issued by the NPPO-(EXPORTING COUNTRY). These pests are currently regulated in Nepal and are restricted to enter into the Nepalese territory.

S.N.	Inherent commodity risk	Additional Declaration	Entry Conditions
1			

Filed by Quarantine and Pesticide Management Centre Special Condition after entry:

1. Product tracking in any suspected condition must answer the following;

- Which quantity of the product is sold to whom
- Proven track of well billing
- Batch or Lot Tracking
- Quality Control

Recommended by

Name of official.....

Post.....

It is hereby agreed that the conditions are accepted by the importer. Post entry Quarantine and inspection in the warehouse is recommended. The importer hereby understands the preconditions set and will obey the conditions accordingly.

.....

Importer's Signature:

FORMAT FOR PRODUCT/COMMODITY RISK ASSESSMENT

Importer is required to upload this document while applying for the EP.

Validity of this report until the Date:

Product name

- 1 Importer's track record
- 2 Exporting country
- 3 Reason(s) why the import is required (For luxury/ for food security/ for medicine/ for breeding/ multiplication)
- 4 Mode(s) of transport
- 5 Status of previous entry condition and follow up study on consumption destinations
- 6 Risk Category (processed/semi processed/raw/finished product for final use)
- 7 Intended processing step after entry in details and the processing destination
- 8 Is the commodity devitalized?
- 9 GMO risk product

10 **Product Identity:**

Provide details on taxonomy and nomenclature

Generally, the taxonomic unit of the product is the species; the use of higher or lower taxonomic units should be supported by a scientifically sound rationale

.....

- 11 National annual production status of product.
- 12 National annual consumption pattern of the product
- 13 Additional Monthly Import Need by volume
- 14 Regulatory status of the product
- 15 Regulatory status of the product elsewhere
- 16 Existing Marketing problem of local product
- 17 What is the level of economic loss to agriculture, forestry or horticulture associated

with this product in its existing geographic range?

Factors to consider

- Reduction effect in local crop yield or quality
- Reduction in local prices or demand, including export markets
- Increase in local production costs (including costs of control)
- Vectors of other pests of economic importance

Rating	Judgement	Confidence Level	Judgement
Non Assessed		Non Assessed	
Negligible		Negligible	
Low		Low	
Medium		Medium	
High		High	

18 What is the level of negative impact on native biodiversity associated with this product in its existing geographic range?

Factors to consider

- Threat to native species, with special focus on threatened and keystone species
- Changed community structure
- Hybridization with native species

Rating	Judgement	Confidence Level	Judgement
Non Assessed		Non Assessed	
Negligible		Negligible	
Low		Low	
Medium		Medium	
High		High	

19 What is the level of negative impact on ecosystem patterns and processes associated with this product in its existing geographic range of Nepal?

Factors to consider

- Physical modifications of habitats
- Changes in nutrient cycling and availability
- Modifications of natural successions
- Changes in trophic and mutualistic interactions

Rating	Judgement	Confidence Level	Judgement
Non Assessed		Non Assessed	
Negligible		Negligible	
Low		Low	
Medium		Medium	
High		High	

20 What is the level of negative social impact associated with this product in its existing geographic range?

Factors to consider

- Unemployment
- Health effects
- Recreation, tourism, education or spiritual impacts
- Aesthetics

Rating	Judgement	Confidence Level	Judgement
Non Assessed		Non Assessed	
Negligible		Negligible	
Low		Low	
Medium		Medium	
High		High	

21 22. What is the level of potential negative impact in the import area (for all sectors) under foreseeable climate change conditions?

Factors to consider

- Climate change projection
- Climate change factors that affect the product’s reproduction and feeding habits
- Number of generations per year

Rating	Judgement	Confidence Level	Judgement
Non Assessed		Non Assessed	
Negligible		Negligible	
Low		Low	
Medium		Medium	
High		High	

Summary of potential consequences

- Summary of rate and confidence level.....
- Summary note on potential consequences.....

Risk Assessment Summary

- Does the product confirm the rational of import?
- What volume by months of the year?
- Any additional condition to be followed by prospective importer
- Additional declaration in the EP?

फाराम नं. ६

शत्रुजिव जोखिम विश्लेषणमा आधारित अन्तरराष्ट्रिय पहुँचको प्रस्तावका लागि निवेदन साथ संलग्न गर्नुपर्ने विवरण

1. निर्यात गर्न चाहेको विरुवा वा विरुवाजन्य वस्तुको परिचय
2. वार्षिक उत्पादनको प्रवृत्ति
3. आन्तरिक खपत
4. उत्पादन प्रकृया
5. मुल्य श्रृंखलाको व्याख्या
6. आपूर्ति श्रृंखलाको व्याख्या
7. वस्तुको गुणस्तर तथा गुणस्तर नियन्त्रणको तरिका
8. प्रस्तावित वार्षिक निर्यात परिमाण
9. समग्र उद्योगको व्यवसायीक योजना

नोट: यस अनुसार व्यवसायी वा सम्बन्धित संस्थाबाट विवरण प्राप्त भएमा अनुसुचि ३ बमोजिमको विवरणहरु समेत संलग्न राखी तयार गरिएको मार्केट एक्सेस सवमिसन डकुमेन्ट संगठनले सम्बन्धित मुलुकमा पठाइ अन्तिम निष्कर्ष नआएसम्म निरन्तर जोखिम संचार गर्ने छ।

